



壁報論文作品欣賞

診所組



The analysis of Diode laser application in facilitated blood coagulation

二極體雷射於促進血液凝結之分析

Abstract

維思登牙醫診所 - 張朝富 醫師


After trauma or injuries, the wounded sites usually result to bleed. The blood will then undergo a physiological process called coagulation to achieve hemostasis in order to prevent excessive blood loss and to make the wound ready for further healing. However, the time required by the blood to start coagulating is determined by the individual's phenotype through its coagulation cascade. There are several known diagnostic diseases or disorders which are related to the impairment of the coagulation cascade such as hemophilia, Von Willebrand disease, and deficiency of vitamin K... etc. In addition, drug application can also be a result of prolonged blood coagulation. This study; however, introduces a clinical method of directly and externally making a pool of blood instantly coagulate by the treatment of Diode laser. For example: to stop bleeding in a dental extraction sockets.

身體因外傷流血時，血液循著凝血連鎖反應機制自然達到凝血作用以避免血液過度流失的情況產生。在某些情況下，例如血友病、von Willebrand 疾病、及維生素 K 的缺乏或是某些藥物的應用，會導致凝血連鎖反應機制受到干擾而不能正常地實行凝血作用。原則上，凝血時間的長短會因人而異，且目前已知鮮少有 " 直接 " 及 " 體外 " 的醫學處置能 " 達到縮短凝血時間的目的。此實驗介紹一種使用牙科二極體雷射透過直接及體外的照射，達到於傷口立即凝血之目的。

LASERS (Light Amplification by Stimulated Emission of Radiation) emit a precise beam of monochromatic, coherent, and collimated light energy. The diode laser is a near-infrared continuous wave laser whose wavelength varies from 655nm to 980nm. The wavelength determines the absorption characteristic in biologic tissues. When the laser beam is applied to the tissue, the light energy is transformed into heat and causes a photo-thermal event resulting a temperature rise in the tissue. Different observed effects take place by different tissue temperature achieved. Coagulation begins at over 50°C, with protein denaturation at 60°C. Vaporization of water occurs at over 100°C, with carbonization of tissue at

200°C. A typical laser surgery is achieved by the process of ablation, meaning tissue removal by converting it to a gaseous state or plume. Lasers have also been utilized to induce vascular occlusion and photocoagulation.

雷射, **LASERS** (Light Amplification by Stimulated Emission of Radiation) 是一種單色、同調及集中的光能量。其中二極體雷射是位於波長範圍655nm至980nm之近紅外線雷射。雷射的波長決定著照射時生物組織中的吸收特質；當雷射照射於組織時，光能被轉換成熱產生光熱效應後導致組織升溫。組織溫度於50°C時血液開始凝結，60°C時組織蛋白質開始變性，達到100°C時水分開始蒸發，組織碳化則發生於溫度 200°C 以上。典型的雷射手術是藉由以組織吸收雷射能量後升溫達到氣化程度進行切割或剝離進行。



Introduction

The laser device chosen in this experiment is ezlase 940 with 400µm laser tip, manufactured by Biolase Technology, Inc.. According to the laser wavelength vs tissue absorption graph, diode laser of wavelength 940nm was designed to be better absorbed by hemoglobin, oxyhemoglobin and water than other types of functional lasers which possess different range of wavelengths. As we had already known that the temperature would rise once the tissue start to absorb energy, and the tissue would eventually coagulate once the temperature reach 60 °C proven by the infrared thermography. Several trials of the study had been conducted previously to determine the appropriate power output, and it was shown that power output of 2.25Watts, 2.50Watts and 2.75Watts were the most effective to produce blood clot. If the power output of less than 2.25Watts was used the blood would hardly form a visible clot, and if more than 2.75 Watts was used, the blood would evaporate immediately without clotting.


In spite of having a variety of tips of different diameter to choose from (200µm, 300µm, 400µm), the diameter of 400µm tip was selected to conduct the experiment as it would theoretically produce blood clot of greater surface area which was more effective and less time consuming.

Aim

The study assessed the level of blood coagulation after diode laser treatment through different power settings and at a different time period comparing to the normal coagulation process.

此實驗以相同的總能量，使用不同的雷射功率以不接觸方式照射人血後分析各組別凝血程度。

Materials and Method



Conclusion

Physiologically, the blood start to coagulate when blood temperature reaches over 50°C because of the denaturation of its protein (i.e. blood cells, serum ,etc...). This facilitated coagulation process is different from the natural coagulation cascade where internal and external pathway and activation of huge amount of various coagulation factors are involved. Depending on the individual, the coagulation process normally takes a certain period of time and cannot be accelerated by any means. However when the individual is in certain pathological status, or has certain blood disorders, the time required by the blood to clot is even prolonged. The experiment introduced a way to make an instant blood clot and thus stop bleeding when a pool of blood in a defect already exists (i.e. dental extraction sockets). According to the results of spectrophotometry test, the Optical Density of laser-treated blood sample groups showed a significant difference than the control group, which proves the blood do form a clot after laser treatment. In addition, since the working principle of the diode laser is to heat-up the chromophore of the target tissue or substance, a functional infrared thermography device was used to directly measure the blood temperature at the time when laser treating, thus explaining the reason of blood clotting due to the physiological mechanism : the blood will denature and coagulate when the temperature achieve over 50°C .

Furthermore, the SEM photo provide the real images of the blood samples and visibly show the blood clot components in nanoscale. Although it is demonstrated that a blood clot in a blood pool can be instantly produced by a diode laser at this experiment, further in vivo studies is needed to prove whether this technique can work in the same manner clinically to achieve facilitated stop-bleeding or if this blood clot formation assisted by laser would cause any negative or positive effects at the surrounding wounded tissues.





壁報論文作品欣賞

診所組

ANTIMICROBIAL PHOTODYNAMIC THERAPY FOR PERIODONTAL TREATMENT

Abstract

潔明牙醫診所 - 吳啟明醫師

The patient was suffering from tooth discomfort of tooth 47 for 3 months and was treated at other clinic by conventional approach. Bony defect was noted upon Panorax-ray examination at the distal site of tooth 47 with a 11mm probing depth pocket. Phase contrast microscopy was used to sample the quantity of microorganism at the pocket site.

There was substantial reduction of microorganism in the pocket after a combination of a Er:br YAG laser periodontal treatment and two antimicrobial photodynamic therapy (aPDT) using methylene blue as photosensitizer. (Please see attached Phase contrast microscope photograph Considerable bone regeneration was detected upon X-ray examination at 6 months follow up appointment.



圖 1 3 (Fig.1) : pre-treatment infrabony defect over #47 distal side



圖 1 4 : Phase contrast microscopy pre-treatment

Treatment Procedure



圖 1 5 (Fig. 2) : Scaling and root planning and Er:Br YAG laser finished



圖 1 6 (Fig. 3) : after SRP 2days, photosensitizer injection (methylene blue) to pocket bottom 3 min.s

The analysis of Diode laser application in facilitated blood coagulation

二極體雷射於促進血液凝結之分析

張朝富

Abstract

After trauma or injuries, the wounded sites usually result to bleed. The blood will then undergo a physiological process called coagulation to achieve hemostasis in order to prevent excessive blood loss and to make the wound ready for further healing. However, the time required by the blood to start coagulating is determined by the individual's phenotype through its coagulation cascade.¹ There are several known diagnostic diseases or disorders which are related to the impairment of the coagulation cascade such as hemophilia, Von Willebrand disease, and deficiency of vitamin K, etc. In addition, drug application can also be a result of prolonged blood coagulation. This study, however, introduces a clinical method of directly and externally making a pool of blood instantly coagulate by the treatment of Diode laser. For example: to stop bleeding in a dental extraction sockets.

LASERS (Light Amplification by Stimulated Emission of Radiation) emit a precise beam of monochromatic, coherent, and collimated light energy. The diode laser is a near-infrared continuous wave laser whose wavelength varies from 655nm to 980nm. The wavelength determines the absorption characteristic in biologic tissues.² When the laser beam is applied to the tissue, the light energy is transformed into heat and causes a photo-thermal event resulting a temperature rise in the tissue. Different observed effects take place by different tissue temperature achieved. Coagulation begins at over 50°C, with protein denaturation at 60°C. Vaporization of water occurs at over 100°C, with carbonization of tissue at 200°C.³ A typical laser surgery is achieved by the process of ablation, meaning tissue removal by converting it to a gaseous state or plume. Lasers have also been utilized to induce vascular occlusion and photocoagulation.

Aim

The study assessed the level of blood coagulation after diode laser treatment through different power settings and at a different time period comparing to the normal coagulation process

Introduction

The laser device chosen in this experiment is ezlase 940 with 400µm laser tip, manufactured by Biolase Technology, Inc. According to the laser wavelength vs tissue absorption graph, diode laser of wavelength 940nm was designed to be better absorbed by hemoglobin, oxyhemoglobin and water than other types of functional lasers which possess different range of wavelengths. As we had already known that the temperature would rise once the tissue start to absorb energy, and the tissue would eventually coagulate once the temperature reach 60°C proven by the infrared thermography. Several trials of the study had been conducted previously to determine the appropriate power output, and it was shown that power output of 2.25Watts, 2.50Watts and 2.75Watts were the most effective to produce blood clot. If the power output of less than 2.25Watts was used the blood would hardly form a visible clot, and if more than 2.75 Watts was used, the blood would evaporate immediately without clotting. In spite of having a variety of tips of different diameter to choose from (200µm, 300µm, 400µm), the diameter of 400µm tip was selected to conduct the experiment as it would theoretically produce blood clot of greater surface area which was more effective and less time consuming.

Materials and Method



Conclusion

Physiologically, the blood start to coagulate when blood temperature reaches over 50°C because of the denaturation of its protein (i.e. blood cells, serum, etc.). This facilitated coagulation process is different from the natural coagulation cascade where internal and external pathway and activation of huge amount of various coagulation factors are involved. Depending on the individual, the coagulation process normally takes a certain period of time and cannot be accelerated by any means. However when the individual is in certain pathological status, or has certain blood disorders, the time required by the blood to clot is even prolonged. The experiment introduced a way to make an instant blood clot and thus stop bleeding when a pool of blood in a defect already exists (i.e. dental extraction sockets). According to the results of spectrophotometry test, the Optical Density of laser-treated blood sample groups showed a significant difference than the control group, which proves the blood do form a clot after laser treatment. In addition, since the working principle of the diode laser is to heat-up the chromophore of the target tissue or substance, a functional infrared thermography device was used to directly measure the blood temperature at the time when laser treating, thus explaining the reason of blood clotting due to the physiological mechanism: the blood will denature and coagulate when the temperature achieve over 50°C. Furthermore, the SEM photo provide the real images of the blood samples and visibly show the blood clot components in nanoscale.

Although it is demonstrated that a blood clot in a blood pool can be instantly produced by a diode laser at this experiment, further *in vivo* studies is needed to prove whether this technique can work in the same manner clinically to achieve facilitated stop-bleeding or if this blood clot formation assisted by laser would cause any negative or positive effects at the surrounding wounded tissues.






摘要

身體因外傷流血時，血液循著凝血連鎖反應機制自然達到凝血作用以避免血液過度流失的情況產生。在某些情況下，例如血友病、von Willebrand 疾病、及維生素K的缺乏或是某些藥物的應用，會導致凝血連鎖反應機制受到干擾而不能正常地實行凝血作用。原則上，凝血時間的長短會因人而異，且目前已知鮮少有“直接”及“體外”的醫學處置能達到縮短凝血時間的目的。此實驗介紹一種使用牙科二極體雷射透過直接及體外的照射，達到於傷口立即凝血之目的。

雷射LASERS (Light Amplification by Stimulated Emission of Radiation) 是一種單色、同調及集中的光能量。其中二極體雷射是位於波長範圍655nm至980nm之近紅外線雷射。雷射的波長決定著照射時生物組織中的吸收特質；當雷射照射於組織時，光能被轉換成熱產生光熱效應導致組織升溫。組織溫度於50°C時血液開始凝結，60°C時組織蛋白質開始變性，達到100°C時水分開始蒸發，組織碳化則發生於溫度200°C以上。典型的雷射手術是藉由以組織吸收雷射能量後升溫達到氧化程度進行切割或剝離進行。

目的

此實驗以相同的總能量，使用不同的雷射功率以不接觸方式照射人血後分析各組別凝血程度。






圖 17 (Fig.4) : 660nm diode laser 10sec per point, 6point per tooth

圖 18 : Phase contrast microscopy Post-treatment



圖 19 (Fig.6) : Post OP 6 months, the newly-formed hard tissue in the periodontal infrabony defect over #47 distal side

圖 20 (Fig.5) : Post-OP 3 month

Discussion

Antimicrobial photodynamic therapy(PDT) as an adjunctive treatment in addition to scaling and root planing for the treatment of periodontitis has been shown to be clinically useful. The underlying mechanism of photodisinfection is the targeting and the elimination of the bacteria most responsible for the progression of periodontal disease. In clinical studies it show better outcome in some cases. In the future PDT may be the choice of treatment for patient and dentist before surgical therapy.

Reference

1. Kao RT et. al. Esthetic crown lengthening: appropriate diagnosis for achieving gingival balance. J Calif Dent Assoc. 2008 Mar;36(3):187-91
2. Magid KS et. al. Laser use for esthetic soft tissue modification. Dent Clin North Am. 2007 Apr;51(2):525-45
3. Polansky R et. al. Clinical effectiveness of photodynamic therapy in the treatment of periodontitis. J Clin Periodontol. 2009 Jul;36(7):575-80.
4. Raghavendra M et. al. Photodynamic therapy: a targeted therapy in periodontics. Aust Dent J. 2009 Sep;54 Suppl 1:S102-9

ANTIMICROBIAL PHOTODYNAMIC THERAPY FOR PERIODONTAL TREATMENT


潔明牙醫診所 吳啟明 醫師

ABSTRACT

The patient was suffering from tooth discomfort of tooth 47 for 3 months and was treated at other clinic by conventional approach. Bony defect was noted upon Panorax x-ray examination at the distal site of tooth 47 with a 11mm probing depth pocket. Phase contrast microscopy was used to sample the quantity of microorganism at the pocket site. There was substantial reduction of microorganism in the pocket after a combination of a Er:br YAG laser periodontal treatment and two antimicrobial photodynamic therapy (aPDT) using methylene blue as photosensitizer. (Please see attached Phase contrast microscope photograph. Considerable bone regeneration was detected upon X-ray examination at 6 months follow up appointment.



Fig.1: pre-treatment infrabony defect over #47 distal side



Phase contrast microscopy pre-treatment

TREATMENT PROCEDURE



Fig. 2: Scaling and root planning and Er:Br YAG laser finished



Fig.3: after SRP 2days, photosensitizer injection (methylene blue) to pocket bottom 3 min.s



Fig.4: 660nm diode laser 10sec per point, 6point per tooth



Phase contrast microscopy Post-treatment



Fig.5: Post-OP 3 month



Fig.6: Post OP 6 months, the newly-formed hard tissue in the periodontal infrabony defect over #47 distal side

DISCUSSION

Antimicrobial photodynamic therapy(PDT) as an adjunctive treatment in addition to scaling and root planing for the treatment of periodontitis has been shown to be clinically useful. The underlying mechanism of photodisinfection is the targeting and the elimination of the bacteria most responsible for the progression of periodontal disease. In clinical studies it show better outcome in some cases. In the future PDT may be the choice of treatment for patient and dentist before surgical therapy.

REFERENCE

1. Kao RT et. al. Esthetic crown lengthening: appropriate diagnosis for achieving gingival balance. J Calif Dent Assoc. 2008 Mar;36(3):187-91
2. Magid KS et. al. Laser use for esthetic soft tissue modification. Dent Clin North Am. 2007 Apr;51(2):525-45
3. Polansky R et. al. Clinical effectiveness of photodynamic therapy in the treatment of periodontitis. J Clin Periodontol. 2009 Jul;36(7):575-80.
4. Raghavendra M et. al. Photodynamic therapy: a targeted therapy in periodontics. Aust Dent J. 2009 Sep;54 Suppl 1:S102-9

壁報論文作品欣賞

診所組



Using Complete Denture and Implant-Supported Overdenture for Treatment of Edentulous Patient with Cleft Palate –A Case Report 二以上顎全口義齒及下顎植體支持覆蓋式義齒治療無牙併上顎裂患

者一病例報告

黃韻璇醫師 晨軒牙醫診所
葉立維醫師 當代牙醫診所

摘要

本病例報告針對一位 65 歲上顎前牙區單側性唇顎裂，具有嚴重牙周病問題，舊活動義齒配戴經驗不佳影響咀嚼功能之男性患者，紀錄如何以開窗印模方法製作上顎全口義齒，下顎則是種植四支人工植體，製作植體支持覆蓋式義齒。治療完成後，活動義齒穩定度提高，病人美觀、發音及咀嚼功能大幅改善。旨針對類似的全口無牙併上顎裂患者之匱復療程提供參考。

病例報告

患者基本資料：

姜先生，65歲，已婚退休男性。沒有食物及藥物過敏史。無抽菸及嚼食檳榔之習慣。

牙科主訴：假牙的咀嚼功能不佳已持續數年，想要重做假牙改善咀嚼功能。

顫顎關節檢查：張口度及張口範圍皆正常，顫顎關節無彈響聲。

口外檢查：

正面觀(frontal view)：臉部有些許不對稱。

微笑觀(smile view)：無微笑露齦(gummy smile)，中線些許不對稱，下顎中線與臉部中線相比往左移1mm。

側面觀(lateral view)：直線輪廓(straight profile)，鼻唇角(nasolabial angle)約90度。
(圖1)

圖1



圖1：患者外觀

口內檢查：

1. 上顎無牙，前牙區齒槽裂 (cleft alveolar ridge) 合併鬆軟組織 (flabby tissue)。唇顎裂形式屬於單側性裂，範圍影響左側唇及齒槽，侷限於原顎 (primary palate) 範圍，即齒槽裂範圍在門齒乳頭 (incisive papilla) 之前，並未延伸到後方的硬顎及軟顎 (secondary palate)。齒槽裂部分由口腔延伸至鼻腔底部，以軟組織相隔，未完全相通。
2. 下顎尚存牙齒為左下正中門齒至左下第二小白齒、左下第三大白齒、右下正中門齒、側門齒及第二小白齒。除左下犬齒及第一小白齒外，其餘牙齒有嚴重之齒槽骨喪失 (圖2、3、4)。
3. 患者之舊假牙為上顎全口義齒、下顎Kennedy Class I-Modification I 之可撤式局部義齒。假牙有嚴重磨耗現象且上顎全口義齒的邊緣過度延伸 (overextension)，上顎前牙區的假牙基底不密合。

圖2



圖3




圖4



圖2：初診之口內影像
圖3：下顎周檢查紀錄表
圖4：初診X光片

治療計畫：

與患者討論之治療計畫有二方案：治療計畫一為上顎全口義齒，左下犬齒及第一小白齒析量牙冠，下顎 Kennedy Class I 可撤式局部義齒。治療計畫二為上顎全口義齒，下顎牙齒全數拔除，種植四支人工植體再予以製作以植體支持之覆蓋式義齒 (implant-supported overdenture)。經與患者討論後，患者選擇治療計畫二。治療流程如圖5至圖10。

治療後追蹤：

治療完成後，活動義齒穩定度提高，病人對於新製作的活動假牙適應良好，美觀、發音及咀嚼功能大幅改善。




圖 5：上顎最終印模 (final impression)
圖 6：下顎最終印模
圖 7：下顎義齒之組織面
圖 8：初診之口內照
圖 9：治療前後之比較

討論

一、本病例上顎印模困難之處：

- (1) 齒槽裂的開口恰位於上顎全口假牙的邊緣 (border) · 邊緣成形 (border molding) 時 · 邊緣的建立要兼顧能達到該交界 (communication) 的邊緣封閉 (seal) 又不能過度延伸 (overextension)¹。
- (2) 齒槽裂處無齒槽骨支撐 · 軟組織似鬆軟組織 (flabby tissue) · 經由文獻回顧對鬆

軟組織的印模後 · 決定以開窗印模技術印模^{2,3}。

二、治療計畫設計方面 · 病人全口牙齒經評估後 · 左下犬齒及第一小白齒是可以考慮留下的。然而考慮患者年紀為65歲 · 左下犬齒及第一小白齒之齒槽骨支持 (alveolar bone support) 約剩百分之五十 · 10年後可能無法留存 · 70幾歲要補種第三象限的植牙風險更大 · 且齒槽骨條件更差。因病人經濟能力許可 · 且根據研究顯示以植體來支持下顎覆蓋式義齒效果較好 · 能有效增加活動假牙的固持力與穩定度 · 並且提高患者的滿意度 · 其植牙的成功率介於97%到100%之間 · 追蹤時間最長可達13年之久^{4,5,6}。有關下顎覆蓋式義齒的文獻 · 多數是使用四支植體支持 · 且植牙成功率可提高至97%⁷。故與患者討論後決定以四支植體完成植體支持覆蓋式義齒。

結論

治療全口無牙併上顎裂患者 · 上顎利用開窗印模技術 · 下顎利用植體支持 · 可改善活動義齒之穩定度 · 進而重建病人美觀 · 發音及咀嚼功能 · 提供此類型患者良好的生活品質。

參考文獻

1. Reisberg DJ. Prosthetic habilitation of Patients with clefts. Clin Plast Surg 2004; 31: 353-60
2. Hrizdana Hadjieva, Mariana Dimova et al. Selective Pressure and Hypermobility Mucosa On The Alveolar Ridges
3. C. D. Lynch¹ and P. F. Allen². Management of the flabby ridge: using contemporary materials to solve an old problem. British Dental Journal 2006; 200: 258-261
4. Koller, B., W. Att, and J. R. Strub. Survival rates of teeth, implants, and double crown-retained removable dental prostheses: a systematic literature review. Int J Prosthodont, 2011. 24(2): p. 109-17
5. Van Steenberghe, D., et al., A prospective evaluation of the fate of 697 consecutive intraoral fixtures and modum Branemark in the rehabilitation of edentulism. J Head Neck Pathol, 1987. 6: p. 53-58.
6. Visser, A., et al., Mandibular overdentures supported by two or four endosseous implants. Clinical oral implants research, 2005. 16(1): p. 19-25
7. ElKerdawy, M. W. and I. A. W. Radi, Effect of dislodging forces on mandibular implant attachment-retained overdenture. Implant dentistry, 2011. 20(3): p. 246-254



壁報論文作品欣賞

診所組

The Influence of Dentist's Experiences On The Behavior of Preschool Children

Huei Wen Liu^{1,2,3,4}, Chen-Yu Hsieh^{2,3,4}, Yen-Kuang Lin³, Chia-Lan Hsiang^{2,3,4}

¹) 葡萄藤兒童牙科, ²) Department of Pediatric Dentistry of Taipei Municipal Wan-Fang Hospital, ³)

Taipei Medical University, ⁴) Taiwan Academy of Pediatric Dentistry

葡萄藤兒童牙科 - 劉惠文醫師

Background & Aim

Behavior management is a key factor in providing dental care for children. Young dentists usually take time to learn the behavior management techniques by trial and error.

- ◆ This study aimed to analyze the behavior management skills of dentists at Taipei Municipal Wan-Fang Hospital, it's influence to the preschool children's behavior at the first visit has also been evaluated.

Design

- ◆ Children aged 3 to 6 years old.
- ◆ First visit at pediatric dentistry department of Taipei Municipal Wan-Fang Hospital were included.
- ◆ All subjects were healthy and denied previous dental experience.
- ◆ Behavior shaping process in the visit was video recorded.
- ◆ The cooperation of patients and the behavior management techniques used by the dentist in the visit were recorded by the author and analyzed by SPSS vol.19.

Results

- ◆ A total of 92 patients were included(50 boys and 42 girls.)
- ◆ There was a positive correlation between the age of subjects and their cooperation during the visit.
- ◆ The pedo resident group treated patients whose age were significantly younger than those in the intern group and the GP resident group.(Figure 1) However, patients in the pedo resident group were significantly more cooperative than the other 2 groups.(Figure 2)



以上顎全口義齒及下顎植體支持覆蓋式義齒治療無牙併上顎裂患者 —病例報告

Using Complete Denture and Implant-Supported Overdenture for Treatment of Edentulous Patient with Cleft Palate – A Case Report

黃韻璇 Yun-Hsuan Huang¹ 葉立維 Li-Wei Yeh²

¹晨軒牙醫診所 Chen Xuan Dental Clinic ²當代牙醫診所 Modern Dental Clinic

摘要

本病例報告針對一位65歲上顎前牙區單側性唇顎裂，具有嚴重牙周病問題，舊活動義齒配戴經驗不佳影響咀嚼功能之男性患者，紀錄如何以閉塞印模方法製作上顎全口義齒，下顎則是種植四支人工植體，製作植體支持覆蓋式義齒。治療完成後，活動義齒穩定度提高，病人美觀、發音及咀嚼功能大幅改善。旨針對類似的全口無牙併上顎裂患者之治療提供參考。

病例報告

患者基本資料：

姜先生，65歲，已婚退休男性，沒有食物及藥物過敏史。無抽菸及嗜食檳榔之習慣。牙科主訴：假牙的咀嚼功能不佳已持續數年，想要重做假牙改善咀嚼功能。

顯微鏡檢查：張口度及張口範圍正常，顯微鏡下無彈簧。

口外檢查：

正面觀(frontal view)：面部有齒許不對稱。

微笑觀(smile view)：無假牙露齒(gummy smile)，中線齒許不對稱。下顎中線與面部中線相比往左移1mm。

側面觀(lateral view)：直線輪廓(straight profile)，鼻唇角(nasolabial angle)

約90度。(圖1)

口內檢查：

1. 上顎無牙，系牙齦齒槽裂(cleft alveolar ridge)合併鬆軟組織(flabby tissue)，唇

顎裂形式屬於單側性裂，範圍影響左側鼻及齒槽，局限於原顎(primary

palate)範圍，即齒槽裂前附在門齒乳頭(incisive papilla)之前，並未延伸到後

方的硬顎及軟顎(secondary palate)。齒槽裂部分由口腔延伸至鼻腔底部，以

後組織相連，未完全相連。

2. 下顎無牙，前牙為左下正中門齒至左下第二小白齒，左下第三大白齒，右下正

中門齒，側門齒及第二小白齒，除左下犬齒及第一小白齒外，其餘牙齒有嚴

重之齒槽骨吸收(圖2、3、4)。

3. 患者之舊假牙為上顎全口義齒，下顎Kennedy Class I 可動式局部義齒。治療計畫二為上

顎全口義齒，下顎牙齒全數拔除，種植四支人工植體再予以製作以植體支持之覆

蓋式義齒(implant-supported overdenture)。經與患者討論後，患者選擇治療

計畫二。治療流程如圖5至圖10。

上顎前牙區的假牙基座不密合。

治療計畫：

與患者討論之治療計畫有二方案：治療計畫一為上顎全口義齒，左下犬齒及第

一小白齒折量牙冠。下顎Kennedy Class I 可動式局部義齒。治療計畫二為上

顎全口義齒，下顎牙齒全數拔除，種植四支人工植體再予以製作以植體支持之覆

蓋式義齒(implant-supported overdenture)。經與患者討論後，患者選擇治療

計畫二。治療流程如圖5至圖10。

治療後追蹤：

治療完成後，活動義齒穩定度提高，病人對於所製作的活動假牙適應良好，美觀、發音及

咀嚼功能大幅改善。



圖1 治療前之臨床照片

圖2 上顎全口義齒配戴前之X光片

圖3 下顎全口義齒配戴前之X光片

圖4 上顎全口義齒配戴前之X光片

圖5 治療計畫二之X光片

圖6 治療計畫二之X光片

圖7 治療計畫二之X光片

圖8 治療計畫二之X光片

圖9 治療計畫二之X光片

圖10 治療計畫二之X光片

圖11 治療計畫二之X光片

圖12 治療計畫二之X光片

圖13 治療計畫二之X光片

圖14 治療計畫二之X光片

圖15 治療計畫二之X光片

圖16 治療計畫二之X光片

圖17 治療計畫二之X光片

圖18 治療計畫二之X光片

圖19 治療計畫二之X光片

圖20 治療計畫二之X光片

圖21 治療計畫二之X光片

圖22 治療計畫二之X光片

圖23 治療計畫二之X光片

圖24 治療計畫二之X光片

圖25 治療計畫二之X光片

圖26 治療計畫二之X光片

圖27 治療計畫二之X光片

圖28 治療計畫二之X光片

圖29 治療計畫二之X光片

圖30 治療計畫二之X光片

圖31 治療計畫二之X光片

圖32 治療計畫二之X光片

圖33 治療計畫二之X光片

圖34 治療計畫二之X光片

圖35 治療計畫二之X光片

圖36 治療計畫二之X光片

圖37 治療計畫二之X光片

圖38 治療計畫二之X光片

圖39 治療計畫二之X光片

圖40 治療計畫二之X光片

圖41 治療計畫二之X光片

圖42 治療計畫二之X光片

圖43 治療計畫二之X光片

圖44 治療計畫二之X光片

圖45 治療計畫二之X光片

圖46 治療計畫二之X光片

圖47 治療計畫二之X光片

圖48 治療計畫二之X光片

圖49 治療計畫二之X光片

圖50 治療計畫二之X光片

圖51 治療計畫二之X光片

圖52 治療計畫二之X光片

圖53 治療計畫二之X光片

圖54 治療計畫二之X光片

圖55 治療計畫二之X光片

圖56 治療計畫二之X光片

圖57 治療計畫二之X光片

圖58 治療計畫二之X光片

圖59 治療計畫二之X光片

圖60 治療計畫二之X光片

圖61 治療計畫二之X光片

圖62 治療計畫二之X光片

圖63 治療計畫二之X光片

圖64 治療計畫二之X光片

圖65 治療計畫二之X光片

圖66 治療計畫二之X光片

圖67 治療計畫二之X光片

圖68 治療計畫二之X光片

圖69 治療計畫二之X光片

圖70 治療計畫二之X光片

圖71 治療計畫二之X光片

圖72 治療計畫二之X光片

圖73 治療計畫二之X光片

圖74 治療計畫二之X光片

圖75 治療計畫二之X光片

圖76 治療計畫二之X光片

圖77 治療計畫二之X光片

圖78 治療計畫二之X光片

圖79 治療計畫二之X光片

圖80 治療計畫二之X光片

圖81 治療計畫二之X光片

圖82 治療計畫二之X光片

圖83 治療計畫二之X光片

圖84 治療計畫二之X光片

圖85 治療計畫二之X光片

圖86 治療計畫二之X光片

圖87 治療計畫二之X光片

圖88 治療計畫二之X光片

圖89 治療計畫二之X光片

圖90 治療計畫二之X光片

圖91 治療計畫二之X光片

圖92 治療計畫二之X光片

圖93 治療計畫二之X光片

圖94 治療計畫二之X光片

圖95 治療計畫二之X光片

圖96 治療計畫二之X光片

圖97 治療計畫二之X光片

圖98 治療計畫二之X光片

圖99 治療計畫二之X光片

圖100 治療計畫二之X光片

圖101 治療計畫二之X光片

圖102 治療計畫二之X光片

圖103 治療計畫二之X光片

圖104 治療計畫二之X光片

圖105 治療計畫二之X光片

圖106 治療計畫二之X光片

圖107 治療計畫二之X光片

圖108 治療計畫二之X光片

圖109 治療計畫二之X光片

圖110 治療計畫二之X光片

圖111 治療計畫二之X光片

圖112 治療計畫二之X光片

圖113 治療計畫二之X光片

圖114 治療計畫二之X光片

圖115 治療計畫二之X光片

圖116 治療計畫二之X光片

圖117 治療計畫二之X光片

圖118 治療計畫二之X光片

圖119 治療計畫二之X光片

圖120 治療計畫二之X光片

圖121 治療計畫二之X光片

圖122 治療計畫二之X光片

圖123 治療計畫二之X光片

圖124 治療計畫二之X光片

圖125 治療計畫二之X光片

圖126 治療計畫二之X光片

圖127 治療計畫二之X光片

圖128 治療計畫二之X光片

圖129 治療計畫二之X光片

圖130 治療計畫二之X光片

圖131 治療計畫二之X光片

圖132 治療計畫二之X光片

圖133 治療計畫二之X光片

圖134 治療計畫二之X光片

圖135 治療計畫二之X光片

圖136 治療計畫二之X光片

圖137 治療計畫二之X光片

圖138 治療計畫二之X光片

圖139 治療計畫二之X光片

圖140 治療計畫二之X光片

圖141 治療計畫二之X光片

圖142 治療計畫二之X光片

圖143 治療計畫二之X光片

圖144 治療計畫二之X光片

圖145 治療計畫二之X光片

圖146 治療計畫二之X光片

圖147 治療計畫二之X光片

圖148 治療計畫二之X光片

圖149 治療計畫二之X光片

圖150 治療計畫二之X光片

圖151 治療計畫二之X光片

圖152 治療計畫二之X光片

圖153 治療計畫二之X光片

圖154 治療計畫二之X光片

圖155 治療計畫二之X光片

圖156 治療計畫二之X光片

圖157 治療計畫二之X光片

圖158 治療計畫二之X光片

圖159 治療計畫二之X光片

圖160 治療計畫二之X光片

圖161 治療計畫二之X光片

圖162 治療計畫二之X光片

圖163 治療計畫二之X光片

圖164 治療計畫二之X光片

圖165 治療計畫二之X光片

圖166 治療計畫二之X光片

圖167 治療計畫二之X光片

圖168 治療計畫二之X光片

圖169 治療計畫二之X光片

圖170 治療計畫二之X光片

圖171 治療計畫二之X光片

圖172 治療計畫二之X光片

圖173 治療計畫二之X光片

圖174 治療計畫二之X光片

圖175 治療計畫二之X光片

圖176 治療計畫二之X光片

圖177 治療計畫二之X光片

圖



- ◆ The pedo resident group used significantly more positive reinforcement and instruction than the other 2 groups during treatment. The intern group and the GP resident groups spent more time on Tell-Show-Do and explanation.(Figure 3)






圖 1 0 : Figure 1. age of subject

圖 1 1 : Figure 2. patient cooperation

圖 1 2 : Figure 3. most used behavior management technique in behavior shaping

Results

- ◆ The older the patient, the more cooperative during first dental visit.
- ◆ Pedo residents treat younger patients with better cooperation.
- ◆ Young dentists should use more positive reinforcement and instruction in behavior shaping.



The Influence of Dentist's Experiences On The Behavior of Preschool Children

Huei Wen Liu^{1,2,3,4}, Chen-Yu Hsieh^{2,3,4}, Yen-Kuang Lin³, Chia-Lan Hsiang^{2,3,4}

¹) 葡萄藤兒童牙科, ²) Department of Pediatric Dentistry of Taipei Municipal Wan-Fang Hospital, ³) Taipei Medical University, ⁴) Taiwan Academy of Pediatric Dentistry



Background & Aim

Behavior management is a key factor in providing dental care for children. Young dentists usually take time to learn the behavior management techniques by trial and error.

- ◆ This study aimed to analyze the behavior management skills of dentists at Taipei Municipal Wan-Fang Hospital, it's influence to the preschool children's behavior at the first visit has also been evaluated.

Design

- ◆ Children aged 3 to 6 years old.
- ◆ First visit at pediatric dentistry department of Taipei Municipal Wan-Fang Hospital were included.
- ◆ All subjects were healthy and denied previous dental experience.
- ◆ Behavior shaping process in the visit was video recorded.
- ◆ The cooperation of patients and the behavior management techniques used by the dentist in the visit were recorded by the author and analyzed by SPSS vol.19.

Results

- ◆ A total of 92 patients were included(50 boys and 42 girls.)
- ◆ There was a positive correlation between the age of subjects and their cooperation during the visit.
- ◆ The pedo resident group treated patients whose age were significantly younger than those in the intern group and the GP resident group.(Figure 1) However, patients in the pedo resident group were significantly more cooperative than the other 2 groups.(Figure 2)
- ◆ The pedo resident group used significantly more positive reinforcement and instruction than the other 2 groups during treatment. The intern group and the GP resident groups spent more time on Tell-Show-Do and explanation.(Figure 3)

Conclusions

- ◆ The older the patient, the more cooperative during first dental visit.
- ◆ Pedo residents treat younger patients with better cooperation.
- ◆ Young dentists should use more positive reinforcement and instruction in behavior shaping.

